# ABclonal®

## Caspase-3 Rabbit pAb

Catalog No.: A0214 65 Publications

## **Basic Information**

#### **Observed MW**

35kDa

#### **Calculated MW**

32kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

The protein encoded by this gene is a cysteine-aspartic acid protease that plays a central role in the execution-phase of cell apoptosis. The encoded protein cleaves and inactivates poly(ADP-ribose) polymerase while it cleaves and activates sterol regulatory element binding proteins as well as caspases 6, 7, and 9. This protein itself is processed by caspases 8, 9, and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease.

## **Recommended Dilutions**

WB 1:500 - 1:1000

**IF/ICC** 1:50 - 1:300

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

## **Immunogen Information**

**Gene ID**836

Swiss Prot
P42574

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

CPP32; SCA-1; CPP32B; Caspase-3

## Contact

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using Caspase-3 Rabbit pAb (A0214) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.