

A0262

Leader in Biomolecular Solutions for Life Science



[KO Validated] CDKN2A/p16INK4a Rabbit pAb

Catalog No.: A0262

KO Validated

73 Publications

Basic Information

Observed MW

16kDa/17kDa

Calculated MW

8kDa/11kDa/12kDa/13kDa/16kDa/17kDa

Category

Polyclonal Antibody

Applications

WB,IF,ICC,ELISA

Cross-Reactivity

Human

Background

This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, the E3 ubiquitin-protein ligase MDM2, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene.

Recommended Dilutions

WB 1:2000 - 1:10000**IF/ICC** 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

1029

Swiss Prot

P42771/Q8N726

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-156 of human CDKN2A/p16INK4a (NP_000068.1).

Synonyms

ARF; MLM; P14; P16; P19; CMM2; INK4; MTS1; TP16; CDK4I; CDKN2; INK4A; MTS-1; P14ARF; P19ARF; P16INK4; P16INK4A; P16-INK4A; 4a

Contact

www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from wild type(WT) and CDKN2A/p16INK4a knockout (KO) 293T(KO) cells, using [KO Validated] CDKN2A/p16INK4a Rabbit pAb (A0262) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25μg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Immunofluorescence analysis of HeLa cells using [KO Validated] CDKN2A/p16INK4a Rabbit pAb (A0262) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.