# ABclonal®

# NTF4 Rabbit pAb

Catalog No.: A0912

# **Basic Information**

#### **Observed MW**

35kDa

#### **Calculated MW**

22kDa

#### **Category**

Polyclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

## **Background**

This gene is a member of a family of neurotrophic factors, neurotrophins, that control survival and differentiation of mammalian neurons. The expression of this gene is ubiquitous and less influenced by environmental signals. While knock-outs of other neurotrophins including nerve growth factor, brain-derived neurotrophic factor, and neurotrophin 3 prove lethal during early postnatal development, NTF5-deficient mice only show minor cellular deficits and develop normally to adulthood.

#### **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

Gene ID Swiss Prot 4909 P34130

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 25-210 of human NTF4 (NP\_006170.1).

#### **Synonyms**

NT4; NT5; NT-4; NT-5; NTF5; GLC10; GLC10; NT-4/5; NTF4

### **Contact**

www.abclonal.com


#### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of lysates from HT-1080 cells, using NTF4 Rabbit pAb (A0912) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 5s.