

A0926

Leader in Biomolecular Solutions for Life Science



NSF Rabbit pAb

Catalog No.: A0926 **1 Publications**

Basic Information

Observed MW

80kDa

Calculated MW

83kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC,IP,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

Enables PDZ domain binding activity and ionotropic glutamate receptor binding activity. Involved in intracellular protein transport; positive regulation of protein catabolic process; and positive regulation of receptor recycling. Located in Golgi apparatus; cytosol; and plasma membrane. Implicated in developmental and epileptic encephalopathy.

Recommended Dilutions

WB 1:1000 - 1:5000

IF/ICC 1:50 - 1:200

IP 0.5µg-4µg antibody for
400µg-600µg extracts
of whole cells

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Contact

 www.abclonal.com

Immunogen Information

Gene ID

4905

Swiss Prot

P46459

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 251-400 of human NSF (NP_006169.2).

Synonyms

SKD2; DEE96; SEC18; NSF

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% Sodium azide,50% glycerol,pH7.3.


Validation Data




Western blot analysis of various lysates using NSF Rabbit pAb (A0926) at 1:3000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 1s.



Immunofluorescence analysis of BALB-3T3 cells using NSF Rabbit pAb (A0926) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U-2 OS cells using NSF Rabbit pAb (A0926) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 600 µg extracts of Mouse brain cells using 3 µg NSF antibody (A0926). Western blot was performed from the immunoprecipitate using NSF antibody (A0926) at a dilution of 1:3000.