# ABclonal® www.abclonal.com

# **TPD52 Rabbit pAb**

Catalog No.: A10254 2 Publications

### **Basic Information**

#### **Observed MW**

26kDa

#### **Calculated MW**

24kDa

#### **Category**

Polyclonal Antibody

#### **Applications**

WB,IHC-P,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Enables calcium ion binding activity and protein homodimerization activity. Involved in B cell differentiation. Located in endoplasmic reticulum and perinuclear region of cytoplasm.

# **Recommended Dilutions**

**WB** 1:1000 - 1:2000

**IHC-P** 1:50 - 1:200

# **Immunogen Information**

**Gene ID**7163
Swiss Prot
P55327

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-184 of human TPD52 (NP\_005070.1).

### **Synonyms**

D52; N8L; PC-1; PrLZ; hD52; TPD52

## **Contact**

www.abclonal.com

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using TPD52 Rabbit pAb (A10254) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded Human tonsil using TPD52 Rabbit pAb (A10254) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain using TPD52 Rabbit pAb (A10254) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.