[KO Validated] CDKN2A/p16INK4a Rabbit pAb

Catalog No.: A11058 KO Validated



Basic Information

Observed MW 17kDa

Calculated MW 8kDa/11kDa/12kDa/13kDa/16kDa/17k Da

Category **Polyclonal Antibody**

Applications WB, IP, ELISA

Cross-Reactivity Human

Background

This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, the E3 ubiquitin-protein ligase MDM2, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene.

Recommended Dilutions

WB	1:500 - 1:1000
IP	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

•	www.abclonal.com

Immunogen Information

Gene ID 1029

Swiss Prot P42771/Q8N726

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-156 of human CDKN2A/p16INK4a (NP_000068.1).

Svnonvms

ARF; MLM; P14; P16; P19; CMM2; INK4; MTS1; TP16; CDK4I; CDKN2; INK4A; MTS-1; P14ARF; P19ARF; P16INK4; P16INK4A; P16-INK4A; 4a

Product Information

Source Rabbit

Isotype IgG

Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Validation Data



Immunoprecipitation analysis of 200 µg extracts of 293T cells, using 3 µg CDKN2A/p16INK4a antibody (A11058). Western blot was performed from the immunoprecipitate using CDKN2A/p16INK4a antibody (A11058) at a dilution of 1:1000.