A11141

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

MonoMethyl-Histone H3-K36 Rabbit mAb

Catalog No.: A11141 Recombinant 2 Publications

Basic Information

Observed MW
16kDa

Calculated MW
16kDa

Category
SMab Recombinant Monoclonal Antibody

Applications
WB,ELISA

Cross-Reactivity
Human,Mouse

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB	1:500 - 1:2000
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
8290

Swiss Prot
Q16695

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; MonoMethyl-Histone H3-K36

Contact

www.abclonal.com

Product Information


Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20℃. Avoid freeze / thaw cycles.
Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using MonoMethyl-Histone H3-K36 Rabbit mAb (A11141).
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.