ABclonal Technology®

www.abclonal.com

[KO Validated] RALBP1 Rabbit pAb

Catalog No.: A1140 **KO Validated**

Basic Information

Observed MW
100kDa

Calculated MW
76kDa

Category
Polyclonal Antibody

Applications
WB,ELISA

Cross-Reactivity
Human,Mouse

Background

RALBP1 plays a role in receptor-mediated endocytosis and is a downstream effector of the small GTP-binding protein RAL (see RALA; MIM 179550). Small G proteins, such as RAL, have GDP-bound inactive and GTP-bound active forms, which shift from the inactive to the active state through the action of RALGDS (MIM 601619), which in turn is activated by RAS (see HRAS; MIM 190020) (summary by Feig, 2003 [PubMed 12888294]).

Recommended Dilutions

WB   1:500 - 1:2000

ELISA   Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID   **Swiss Prot**
10928   Q15311

Immunogen
Recombinant fusion protein containing a sequence corresponding to amino acids 396-655 of human RALBP1 (NP_006779.1).

Synonyms
RIP1; RLIP1; RLIP76; P1

Contact

🌐   www.abclonal.com

Product Information

Source   **Isotype**   **Purification**
Rabbit   IgG   Affinity purification

Storage
Store at -20℃. Avoid freeze / thaw cycles.
Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data

Western blot analysis of various lysates using [KO Validated] RALBP1 Rabbit pAb (A1140) at 1:500 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25μg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.

Western blot analysis of lysates from wild type (WT) and RALBP1 knockout (KO) HeLa cells, using [KO Validated] RALBP1 Rabbit pAb (A1140) at 1:500 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25μg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 5s.