A11463

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Histone H2AX Rabbit pAb

Catalog No.: A11463 7 Publications

Basic Information

Observed MW
15kDa

Calculated MW
15kDa

Category
Polyclonal Antibody

Applications
WB,IHC-P,IF/ICC,IP,ELISA

Cross-Reactivity
Human,Mouse,Rat,Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
IP	0.5μg-4μg antibody for 200μg-400μg extracts of whole cells
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
3014

Swiss Prot
P16104

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
H2A.X; H2A/X; H2AFX; Histone H2AX

Product Information

Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃


Contact

www.abclonal.com


Validation Data




Western blot analysis of various lysates using Histone H2AX Rabbit pAb (A11463) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 15s.




Immunohistochemistry analysis of paraffin-embedded Human liver cancer using Histone H2AX Rabbit pAb (A11463) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human placenta using Histone H2AX Rabbit pAb (A11463) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney using Histone H2AX Rabbit pAb (A11463) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver using Histone H2AX Rabbit pAb (A11463) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Confocal immunofluorescence analysis of HeLa cells using Histone H2AX Rabbit pAb (A11463) at dilution of 1:400. Blue: DAPI for nuclear staining.




Confocal immunofluorescence analysis of U-2 OS cells using Histone H2AX Rabbit pAb (A11463) at dilution of 1:400. Blue: DAPI for nuclear staining.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit https://abclonal.com for a complete listing of recommended products.

Validation Data



Immunofluorescence analysis of U2OS cells using Histone H2AX Rabbit pAb (A11463) at dilution of 100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 200 µg extracts of HeLa cells, using 3 µg Histone H2AX antibody (A11463). Western blot was performed from the immunoprecipitate using Histone H2AX antibody (A11463) at a dilution of 1:1000.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit https://abclonal.com for a complete listing of recommended products.