# ABclonal www.abclonal.com

## AIFM2/FSP1 Rabbit pAb

Catalog No.: A12128 6 Publications

## **Basic Information**

#### **Observed MW**

41kDa

#### **Calculated MW**

41kDa

#### Category

Polyclonal Antibody

## **Applications**

WB,IHC-P,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a flavoprotein oxidoreductase that binds single stranded DNA and is thought to contribute to apoptosis in the presence of bacterial and viral DNA. The expression of this gene is also found to be induced by tumor suppressor protein p53 in colon cancer cells.

## **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IHC-P** 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**84883

Swiss Prot
Q9BRQ8

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 43-371 of human AIFM2/FSP1/AMID/AMID (NP\_116186.1).

#### **Synonyms**

AMID; FSP1; PRG3; AIFM2/FSP1

## Contact


www.abclonal.com

## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from HepG2 cells, using AIFM2/FSP1 Rabbit pAb (A12128) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 180s.




Western blot analysis of various lysates using AIFM2/FSP1 Rabbit pAb (A12128) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer using AIFM2/FSP1 Rabbit pAb (A12128) at dilution of 1:20 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human liver cancer using AIFM2/FSP1 Rabbit pAb (A12128) at dilution of 1:20 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.