A12295

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

PIK3C3/VPS34 Rabbit mAb

Catalog No.: A12295 Recombinant 5 Publications

Basic Information

Observed MW
102kDa

Calculated MW
102kDa

Category
Monoclonal Antibody

Applications
WB,IF/ICC,IP,ELISA

Cross-Reactivity
Human,Mouse,Rat

Clone number
ARC0286

Background

Enables 1-phosphatidylinositol-3-kinase activity. Involved in early endosome to late endosome transport and regulation of cytokinesis. Acts upstream of or within autophagy and protein lipidation. Located in autolysosome; late endosome; and midbody.

Recommended Dilutions

WB	1:1000 - 1:6000
IF/ICC	1:200 - 1:800
IP	0.5μg-4μg antibody for 400μg-600μg extracts of whole cells
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
5289

Swiss Prot
Q8NEB9

Immunogen
Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms
VPS34; Vps34; hVps34; PIK3C3/VPS34

Product Information

Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃


Contact

www.abclonal.com


Validation Data




Western blot analysis of various lysates using PIK3C3/VPS34 Rabbit mAb (A12295) at 1 : 1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.



Confocal imaging of PC-3 cells using PIK3C3/VPS34 Rabbit mAb (A12295, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007,dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Confocal imaging of C2C12 cells using PIK3C3/VPS34 Rabbit mAb (A12295, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunoprecipitation analysis of 600 µg extracts of Rat brain cells using 3 µg PIK3C3/VPS34 antibody (A12295). Western blot was performed from the immunoprecipitate using PIK3C3/VPS34 antibody (A12295) at a dilution of 1:500.