# **PCNA Rabbit mAb**

Catalog No.: A12427 Recombinant 55 Publications



## **Basic Information**

#### **Observed MW**

36kDa

### **Calculated MW**

29kDa

### Category

Monoclonal Antibody

### **Applications**

WB,IHC-P,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat, African green monkey

### CloneNo number

ARC51324

# **Background**

The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome.

# **Recommended Dilutions**

**WB** 1:5000 - 1:20000

IHC-P 1:10000 - 1:40000

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
5111
P12004

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

ATLD2; PCNA

## **Contact**

www.abclonal.com

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# Validation Data



Western blot analysis of various lysates using PCNA Rabbit mAb (A12427) at 1:5000 dilution incubated overnight at  $4^{\circ}$ C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.