SLC12A1 Rabbit pAb

Catalog No.: A12999 2 Publications



Basic Information

Observed MW 150kDa

Calculated MW 121kDa

Category **Polyclonal Antibody**

Applications WB, ELISA

Cross-Reactivity Mouse,Rat

Background

This gene encodes a kidney-specific sodium-potassium-chloride cotransporter that is expressed on the luminal membrane of renal epithelial cells of the thick ascending limb of Henle's loop and the macula densa. It plays a key role in concentrating urine and accounts for most of the NaCl resorption. It is sensitive to such diuretics as furosemide and bumetanide. Some Bartter-like syndromes result from defects in this gene. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional splice variants have been described but their biological validity in humans has not been experimentally proven.

Recommended Dilutions Immunogen Information

WB

1:500 - 1:2000

Gene ID 6557

Swiss Prot Q13621

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 50-150 of human SLC12A1 (NP_000329.2).

Synonyms

BSC; BSC1; CCC2; BSC-1; NKCC2; SLC12A1

Contact

Product Information

www.abclonal.com G

Source

Isotype IgG

Purification Affinity purification

Storage

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using SLC12A1 Rabbit pAb (A12999) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.