# ABclonal www.abclonal.com

# DCP1A Rabbit pAb

Catalog No.: A13431

#### **Basic Information**

#### **Observed MW**

60kDa

#### **Calculated MW**

63kDa

#### Category

Polyclonal Antibody

### **Applications**

WB

#### **Cross-Reactivity**

Human, Mouse

# **Background**

Decapping is a key step in general and regulated mRNA decay. The protein encoded by this gene is a decapping enzyme. This protein and another decapping enzyme form a decapping complex, which interacts with the nonsense-mediated decay factor hUpf1 and may be recruited to mRNAs containing premature termination codons. This protein also participates in the TGF-beta signaling pathway. Alternative splicing of this gene results in multiple transcript variants.

#### **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 

**Swiss Prot** 

55802

Q9NPI6

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

SMIF; SMAD4IP1; HSA275986; Nbla00360; DCP1A

## Contact

€

www.abclonal.com

#### **Product Information**

Source

Isotype

**Purification** 

Rabbit

IgG


Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of lysates from HeLa cells, using DCP1A Rabbit pAb (A13431). Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.