# ABclonal www.abclonal.com

## **IRF8 Rabbit pAb**

Catalog No.: A13677

## **Basic Information**

#### **Observed MW**

50kDa

#### **Calculated MW**

48kDa

## **Category**

Polyclonal Antibody

#### **Applications**

WB, ELISA

## **Cross-Reactivity**

Human

## **Background**

Interferon consensus sequence-binding protein (ICSBP) is a transcription factor of the interferon (IFN) regulatory factor (IRF) family. Proteins of this family are composed of a conserved DNA-binding domain in the N-terminal region and a divergent C-terminal region that serves as the regulatory domain. The IRF family proteins bind to the IFN-stimulated response element (ISRE) and regulate expression of genes stimulated by type I IFNs, namely IFN-alpha and IFN-beta. IRF family proteins also control expression of IFN-alpha and IFN-beta-regulated genes that are induced by viral infection.

## **Recommended Dilutions**

WB 1:100 - 1:500

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**3394

Swiss Prot
Q02556

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

ICSBP; IRF-8; ICSBP1; IMD32A; IMD32B; H-ICSBP; IRF8

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of lysates from HL-60 cells, using IRF8 Rabbit pAb (A13677) at 1:400 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.