ACAT2 Rabbit mAb

Catalog No.: A1399 Recombinant 3 Publications

Background

this gene.



Basic Information

Observed MW 41kDa

Calculated MW 41kDa

Category SMab Recombinant Monoclonal Antibody

Applications WB, ELISA

Cross-Reactivity Human, Mouse, Rat

CloneNo number ARC2560

Recommended Dilutions Immunogen Information

WB

1:500 - 1:1000

Gene ID 39

Swiss Prot Q9BWD1

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 150-250 of human ACAT2 (Q9BWD1).

The product of this gene is an enzyme involved in lipid metabolism, and it encodes

cytosolic acetoacetyl-CoA thiolase. This gene shows complementary overlapping with the 3-prime region of the TCP1 gene in both mouse and human. These genes are encoded on opposite strands of DNA, as well as in opposite transcriptional orientation.

Alternatively spliced transcript variants encoding different isoforms have been found for

Synonyms

ACAT2; acetyl-CoA acetyltransferase 2

Contact	
Contact	

Product Information

€ www.abclonal.com

Isotype IgG

Purification Affinity purification

Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using ACAT2 antibody (A1399) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Western blot analysis of extracts of various cell lines, using ACAT2 antibody (A1399) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.