# LPCAT4 Rabbit pAb

Catalog No.: A14316



#### **Basic Information**

**Observed MW** 57kDa

**Calculated MW** 57kDa

Category **Polyclonal Antibody** 

Applications WB,ELISA

**Cross-Reactivity** Human, Mouse

## Background

Members of the 1-acylglycerol-3-phosphate O-acyltransferase (EC 2.3.1.51) family, such as AGPAT7, catalyze the conversion of lysophosphatidic acid (LPA) to phosphatidic acid (PA), a precursor in the biosynthesis of all glycerolipids. Both LPA and PA are involved in signal transduction (Ye et al., 2005 [PubMed 16243729]).

## **Recommended Dilutions**

WB	1:500 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

Gene ID	Swiss Prot
254531	Q643R3
Immunogen	

Recombinant fusion protein containing a sequence corresponding to amino acids 345-524 of human LPCAT4 (NP\_705841.2).

#### **Synonyms**

AYTL3; AGPAT7; LPEAT2; LPLAT10; LPAAT-eta; LPCAT4

## **Product Information**

Ð www.abclonal.com


Source Rabbit

Isotype IgG

Purification Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.



Western blot analysis of various lysates using LPCAT4 Rabbit pAb (A14316) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.