# DAD1 Rabbit pAb

Catalog No.: A14723 2 Publications



### **Basic Information**

Observed MW 22kDa

Calculated MW 12kDa

**Category** Polyclonal Antibody

Applications WB,ELISA

**Cross-Reactivity** Human, Mouse, Rat

### Background

DAD1, the defender against apoptotic cell death, was initially identified as a negative regulator of programmed cell death in the temperature sensitive tsBN7 cell line. The DAD1 protein disappeared in temperature-sensitive cells following a shift to the nonpermissive temperature, suggesting that loss of the DAD1 protein triggered apoptosis. DAD1 is believed to be a tightly associated subunit of oligosaccharyltransferase both in the intact membrane and in the purified enzyme, thus reflecting the essential nature of N-linked glycosylation in eukaryotes.

# Recommended Dilutions

WB	1:500 - 1:2000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize
	the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID** 1603 Swiss Prot P61803

#### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

# Synonyms

OST2; DAD1

# Contact

# **Product Information**

www.abclonal.com

#### **Source** Rabbit

**Isotype** IgG

**Purification** Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using DAD1 Rabbit pAb (A14723) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5min.