# ABclonal®

# RNF170 Rabbit pAb

Catalog No.: A15195 1 Publications

# **Basic Information**

#### **Observed MW**

30kDa

#### **Calculated MW**

30kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

## **Cross-Reactivity**

Mouse

# **Background**

This gene encodes a RING domain-containing protein that resides in the endoplasmic reticulum (ER) membrane. This protein functions as an E3 ubiquitin ligase and mediates ubiquitination and processing of inositol 1,4,5-trisphosphate (IP3) receptors via the ER-associated protein degradation pathway. It is recruited to the activated IP3 receptors by the ERLIN1/ERLIN2 complex to which it is constitutively bound. Mutations in this gene are associated with autosomal dominant sensory ataxia. Alternatively spliced transcript variants have been found for this gene.

# **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**81790

Swiss Prot
Q96K19

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 46-120 of human RNF170 (NP\_001153696.1).

#### **Synonyms**

ADSA; SNAX1; SPG85; RNF170

## **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using RNF170 Rabbit pAb (A15195) at 1:1000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 15s.