

A1525

Leader in Biomolecular Solutions for Life Science



Pan Acetyl-Lysine Mouse mAb

Catalog No.: A1525

Basic Information

Observed MW

Calculated MW

Category

Monoclonal Antibody

Applications

WB,ELISA

Cross-Reactivity

Human,Mouse,Rat,Other (Wide Range Predicted)

CloneNo number

AMC0491

Background

Acetylation of lysine, like phosphorylation of serine, threonine or tyrosine, is an important reversible modification controlling protein activity. The conserved amino-terminal domains of the four core histones (H2A, H2B, H3, and H4) contain lysines that are acetylated by histone acetyltransferases (HATs) and deacetylated by histone deacetylases (HDACs) (PMID: 9667866). Signaling resulting in acetylation/deacetylation of histones, transcription factors, and other proteins affects a diverse array of cellular processes including chromatin structure and gene activity, cell growth, differentiation, and apoptosis (PMID: 14593721). Recent proteomic surveys suggest that acetylation of lysine residues may be a widespread and important form of post-translational protein modification that affects thousands of proteins involved in control of cell cycle and metabolism, longevity, actin polymerization, and nuclear transport (PMID: 19608861). The regulation of protein acetylation status is impaired in cancer and polyglutamine diseases (PMID: 11864588), and HDACs have become promising targets for anti-cancer drugs currently in development (PMID: 15032670).

Recommended Dilutions

WB 1:500 - 1:1000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Recombinant fusion protein corresponding to a sequence containing acetylated K.

Synonyms

Contact

 www.abclonal.com

Product Information

Source

Mouse

Isotype

IgG1

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using Pan Acetyl-Lysine Mouse mAb (A1525) at 1:1000 dilution. HeLa, NIH/3T3 and C6 cells were treated by TSA (1 uM) at 37°C for 18 hours.

Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.