OXGR1 Rabbit pAb

Catalog No.: A15431



Basic Information

Observed MW 38kDa

Calculated MW 38kDa

Category Polyclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human, Mouse, Rat

Background

This gene encodes a G protein-coupled receptor (GPCR) that belongs to the oxoglutarate receptor family within the GPCR superfamily. The encoded protein is activated by the citric acid intermediate, oxoglutarate, as well as several cysteinyl leukotrienes, including leukotrienes E4, C4 and D4, which are implicated in many inflammatory disorders. In mice, a knock-out of this gene leads to middle ear inflammation, changes in the mucosal epithelium, and an increase in fluid behind the eardrum, and is associated with hearing loss. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB1:200 - 1:2000ELISARecommended starting
concentration is 1
μg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Immunogen Information

Gene ID 27199

Swiss Prot Q96P68

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

aKGR; GPR80; GPR99; P2Y15; P2RY15; OXGR1

Contact

Product Information

www.abclonal.com


Source Rabbit

Isotype IgG **Purification** Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using OXGR1 Rabbit pAb (A15431) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.