Leader in Biomolecular Solutions for Life Science

# **POMK Rabbit pAb**

Catalog No.: A15529



### **Basic Information**

### **Observed MW**

45kDa

#### **Calculated MW**

40kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Mouse,Rat

### **Background**

This gene encodes a protein that may be involved in the presentation of the laminin-binding O-linked carbohydrate chain of alpha-dystroglycan (a-DG), which forms transmembrane linkages between the extracellular matrix and the exoskeleton. Some pathogens use this O-linked carbohydrate unit for host entry. Loss of function compound heterozygous mutations in this gene were found in a human patient affected by the Walker-Warburg syndrome (WWS) phenotype. Mice lacking this gene contain misplaced neurons (heterotopia) in some regions of the brain, possibly from defects in neuronal migration. Alternative splicing of this gene results in multiple transcript variants.

### **Recommended Dilutions**

WB

1:200 - 1:2000

### Immunogen Information

**Gene ID** 84197

Swiss Prot 09H5K3

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-350 of human POMK (NP\_115613.1).

### **Synonyms**

SGK196; MDDGA12; MDDGC12; POMK

#### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using POMK Rabbit pAb (A15529) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.