# ABclonal® www.abclonal.com

## STT3B Rabbit pAb

Catalog No.: A15574

## **Basic Information**

#### **Observed MW**

102kDa

#### **Calculated MW**

94kDa

#### Category

Polyclonal Antibody

## **Applications**

WB,ELISA

## **Cross-Reactivity**

Human, Mouse

## **Background**

The protein encoded by this gene is a catalytic subunit of a protein complex that transfers oligosaccharides onto asparagine residues. Defects in this gene are a cause of congenital disorder of glycosylation Ix (CDG1X).

## **Recommended Dilutions**

**WB** 1:1000 - 1:5000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**201595

Swiss Prot
Q8TCJ2

## **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-100 of human STT3B (NP\_849193.1).

#### **Synonyms**

SIMP; CDG1X; STT3-B; STT3B

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from HUH-7 cells, using STT3B Rabbit pAb (A15574) at 1:3000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.