Leader in Biomolecular Solutions for Life Science

Sorbitol Dehydrogenase Rabbit pAb

www.abclonal.com

ABclonal

Catalog No.: A15727

Basic Information

Observed MW

38kDa

Calculated MW

38kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Sorbitol dehydrogenase (SORD; EC 1.1.1.14) catalyzes the interconversion of polyols and their corresponding ketoses, and together with aldose reductase (ALDR1; MIM 103880), makes up the sorbitol pathway that is believed to play an important role in the development of diabetic complications (summarized by Carr and Markham, 1995 [PubMed 8535074]). The first reaction of the pathway (also called the polyol pathway) is the reduction of glucose to sorbitol by ALDR1 with NADPH as the cofactor. SORD then oxidizes the sorbitol to fructose using NAD(+) cofactor.

Recommended Dilutions

WB

1:500 - 1:2000

Immunogen Information

Gene ID 6652

Swiss Prot Q00796

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-357 of human Sorbitol Dehydrogenase (NP_003095.2).

Synonyms

RDH; SDH; XDH; SORD1; SORDD; HEL-S-95n; Sorbitol Dehydrogenase

Contact

0

www.abclonal.com

Product Information

Source Rabbit

Isotype IgG


Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using Sorbitol Dehydrogenase Rabbit pAb (A15727) at 1:500 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.