# HEY1 Rabbit pAb

Catalog No.: A16110 11 Publications



# **Basic Information**

**Observed MW** 33kDa

**Calculated MW** 33kDa

Category **Polyclonal Antibody** 

Applications WB, IF/ICC, ELISA

**Cross-Reactivity** Human, Mouse, Rat

# Background

This gene encodes a nuclear protein belonging to the hairy and enhancer of splitrelated (HESR) family of basic helix-loop-helix (bHLH)-type transcriptional repressors. Expression of this gene is induced by the Notch and c-Jun signal transduction pathways. Two similar and redundant genes in mouse are required for embryonic cardiovascular development, and are also implicated in neurogenesis and somitogenesis. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

WB	1:500 - 1:1000
IF/ICC	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

Gene ID	Swiss Prot
23462	Q9Y5J3

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 114-129 of human HEY1 (NP\_001269780.1).

#### **Synonyms**

CHF2; OAF1; HERP2; HESR1; HRT-1; NERP2; hHRT1; BHLHb31; HEY1

# Contact

# **Product Information**

Ð www.abclonal.com

Source Rabbit


Isotype IgG

Purification Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using HEY1 Rabbit pAb (A16110) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of A431 cells using HEY1 Rabbit pAb (A16110) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.