

A16218

Leader in Biomolecular Solutions for Life Science



MyoD1 Rabbit pAb

Catalog No.: A16218

4 Publications

Basic Information

Observed MW

45kDa

Calculated MW

35kDa

Category

Polyclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

This gene encodes a nuclear protein that belongs to the basic helix-loop-helix family of transcription factors and the myogenic factors subfamily. It regulates muscle cell differentiation by inducing cell cycle arrest, a prerequisite for myogenic initiation. The protein is also involved in muscle regeneration. It activates its own transcription which may stabilize commitment to myogenesis.

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

4654

Swiss Prot

P15172

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

PUM; MYF3; MYOD; CMYP17; bHLHc1; MYODRIF; MyoD1

Contact

 www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG


Purification

Affinity purification

Storage

Store at -20°C

Validation Data



Western blot analysis of various lysates using MyoD1 Rabbit pAb (A16218) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Western blot analysis of lysates from Rat skeletal muscle, using MyoD1 Rabbit pAb (A16218) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Immunofluorescence analysis of RD cells using MyoD1 Rabbit pAb (A16218) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.