

A17112

Leader in Biomolecular Solutions for Life Science



## PIK3R3/p55PIK Rabbit pAb

Catalog No.: A17112

2 Publications

### Basic Information

#### Observed MW

55kDa

#### Calculated MW

54kDa

#### Category

Polyclonal Antibody

#### Applications

WB,IF/ICC,ELISA

#### Cross-Reactivity

Human,Mouse,Rat

### Background

Phosphatidylinositol 3-kinase (PI3K) phosphorylates phosphatidylinositol and similar compounds, which then serve as second messengers in growth signaling pathways. PI3K is composed of a catalytic and a regulatory subunit. The protein encoded by this gene represents a regulatory subunit of PI3K. The encoded protein contains two SH2 domains through which it binds activated protein tyrosine kinases to regulate their activity.

### Recommended Dilutions

WB 1:100 - 1:500

IF/ICC 1:50 - 1:100

### Immunogen Information

#### Gene ID

8503

#### Swiss Prot

Q92569

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 300-388 of human PIK3R3/p55PIK (NP\_003620.3).

#### Synonyms

p55; p55PIK; p55-GAMMA; PIK3R3/p55PIK

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of various lysates using PIK3R3/p55PIK Rabbit pAb (A17112) at 1:500 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of NIH-3T3 cells using PIK3R3/p55PIK Rabbit pAb (A17112) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.