A17465

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

# CRY2 Rabbit pAb

**Catalog No.: A17465** 3 Publications

## Basic Information

**Observed MW**
70kDa/

**Calculated MW**
67kDa

**Category**
Polyclonal Antibody

**Applications**
WB,ELISA

**Cross-Reactivity**
Human,Mouse,Rat

## Background

This gene encodes a flavin adenine dinucleotide-binding protein that is a key component of the circadian core oscillator complex, which regulates the circadian clock. This gene is upregulated by CLOCK/ARNTL heterodimers but then represses this upregulation in a feedback loop using PER/CRY heterodimers to interact with CLOCK/ARNTL. Polymorphisms in this gene have been associated with altered sleep patterns. The encoded protein is widely conserved across plants and animals. Two transcript variants encoding different isoforms have been found for this gene.

## Recommended Dilutions

| | |
|---|---|
| **WB** | 1:500 - 1:1000 |
| **ELISA** | Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements. |

## Immunogen Information

**Gene ID**
1408

**Swiss Prot**
Q49AN0

**Immunogen**
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

**Synonyms**
HCRY2; PHLL2; CRY2

## Contact

www.abclonal.com

## Product Information


**Source**
Rabbit

**Isotype**
IgG


**Purification**
Affinity purification

**Storage**
Store at -20℃

## Validation Data




Western blot analysis of various lysates, using CRY2 Rabbit pAb (A17465) at 1:400 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit (RM00021).
Exposure time: 180s.




Western blot analysis of various lysates using CRY2 Rabbit pAb (A17465) at 1:400 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates / proteins: 25 µg per lane.
Blocking buffer: 3 % nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.