K11-linkage Specific Polyubiquitin Rabbit pAb

Catalog No.: A18197 1 Publications



Basic Information

Observed MW

Calculated MW

Category Polyclonal Antibody

Applications WB,ELISA,DB

Cross-Reactivity Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Polyubiquitination is a posttranslational modification where ubiquitin chains containing isopeptide bonds linking one of seven ubiquitin lysines with the C terminus of an adjoining ubiquitin are covalently attached to proteins. A crystal structure of K11-linked diubiquitin demonstrates a distinct conformation from K48- or K63-linked diubiquitin. It is important that K11-linked ubiquitin chains as critical regulators of mitotic protein degradation.

Recommended Dilutions

WB	1:500 - 1:2000
DB	1:500 - 1:1000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

1:500 - 1:2000	Gene ID	Swiss Prot
1:500 - 1:1000	Immunogen Synthetic peptide. This	information is considered to be commercially sensitive.
Recommended starting concentration is 1		
the concentration is 1 µg/mL. Please optimize the concentration based on your specific	Synonyms	
assay requirements.		

Contact	

Product Information

€ www.abclonal.com Source Rabbit


Isotype IgG

Purification Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% Sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using K11-linkage Specific Polyubiquitin Rabbit pAb (A18197) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 90s.



Dot-blot analysis of all sorts of peptides using K11-linkage Specific Polyubiquitin antibody (A18197) at 1:1000 dilution.