# ABclonal® www.abclonal.com

## **DEFA5 Rabbit pAb**

Catalog No.: A18208 2 Publications

## **Basic Information**

## **Observed MW**

12kDa

#### **Calculated MW**

10kDa

#### **Category**

Polyclonal Antibody

#### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human

## **Background**

Defensins are a family of antimicrobial and cytotoxic peptides thought to be involved in host defense. They are abundant in the granules of neutrophils and also found in the epithelia of mucosal surfaces such as those of the intestine, respiratory tract, urinary tract, and vagina. Members of the defensin family are highly similar in protein sequence and distinguished by a conserved cysteine motif. Several of the alpha defensin genes appear to be clustered on chromosome 8. The protein encoded by this gene, defensin, alpha 5, is highly expressed in the secretory granules of Paneth cells of the ileum.

## **Recommended Dilutions**

**WB** 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
Q01523

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 20-94 of human DEFA5 (NP\_066290.1).

#### **Synonyms**

DEF5; HD-5; DEFA5

## Contact

www.abclonal.com


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from HCT116 cells, using DEFA5 Rabbit pAb (A18208) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.