# ABclonal www.abclonal.com

# MRPL55 Rabbit pAb

Catalog No.: A18242

# **Basic Information**

#### **Observed MW**

15kDa

#### **Calculated MW**

15kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

# **Cross-Reactivity**

Human

# **Background**

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 39S subunit protein. Multiple transcript variants encoding two different isoforms were identified through sequence analysis.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 

**Swiss Prot** 

128308

Q7Z7F7

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-128 of human MRPL55 (NP\_001308213.1).

### **Synonyms**

L55nt; MRP-L55; AAVG5835; PRO19675; MRPL55

# **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of lysates from MCF7 cells, using MRPL55 Rabbit pAb (A18242) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.