A18257

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

[KO Validated] ATM Rabbit pAb

Catalog No.: A18257 KO Validated 2 Publications

Basic Information

Observed MW
350kDa

Calculated MW
351kDa

Category
Polyclonal Antibody

Applications
WB,ELISA

Cross-Reactivity
Human

Background

The protein encoded by this gene belongs to the PI3/PI4-kinase family. This protein is an important cell cycle checkpoint kinase that phosphorylates; thus, it functions as a regulator of a wide variety of downstream proteins, including tumor suppressor proteins p53 and BRCA1, checkpoint kinase CHK2, checkpoint proteins RAD17 and RAD9, and DNA repair protein NBS1. This protein and the closely related kinase ATR are thought to be master controllers of cell cycle checkpoint signaling pathways that are required for cell response to DNA damage and for genome stability. Mutations in this gene are associated with ataxia telangiectasia, an autosomal recessive disorder.

Recommended Dilutions

WB	1:500 - 1:1000
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
472

Swiss Prot
Q13315

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
AT1; ATA; ATC; ATD; ATE; ATDC; TEL1; TELO1; TM

Contact

www.abclonal.com

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃

Validation Data



Western blot analysis of various lysates using [KO Validated] ATM Rabbit pAb (A18257) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 3min.



Western blot analysis of lysates from wild type (WT) and ATM knockout (KO) 293T cells, using [KO Validated] ATM Rabbit pAb (A18257) at 1:500 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Enhanced Kit (RM00021).
Exposure time: 180s.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit https://abclonal.com for a complete listing of recommended products.