# ABclonal www.abclonal.com

# [KO Validated] Rad23A Rabbit pAb

Catalog No.: A19884

**KO** Validated 1 Publications

### **Basic Information**

#### **Observed MW**

49kDa

### **Calculated MW**

40kDa

### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse

## **Background**

The protein encoded by this gene is one of two human homologs of Saccharomyces cerevisiae Rad23, a protein involved in nucleotide excision repair. Proteins in this family have a modular domain structure consisting of an ubiquitin-like domain (UbL), ubiquitin-associated domain 1 (UbA1), XPC-binding domain and UbA2. The protein encoded by this gene plays an important role in nucleotide excision repair and also in delivery of polyubiquitinated proteins to the proteasome. Alternative splicing results in multiple transcript variants encoding multiple isoforms.

# **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

### **Immunogen Information**

**Gene ID Swiss Prot** 5886 P54725

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

HR23A; HHR23A; 3A

### **Contact**

www.abclonal.com €


### **Product Information**

Source **Isotype Purification** Rabbit Affinity purification IgG

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from wild type (WT) and Rad23A knockout (KO) 293T cells, using [KO Validated] Rad23A Rabbit pAb (A19884) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.