

# [KO Validated] Integrin beta 4 (ITGB4/CD104) Rabbit pAb

Catalog No.: A19890 KO Validated

## **Basic Information**

#### **Observed MW**

230kDa

#### **Calculated MW**

202kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

Integrins are heterodimers comprised of alpha and beta subunits, that are noncovalently associated transmembrane glycoprotein receptors. Different combinations of alpha and beta polypeptides form complexes that vary in their ligand-binding specificities. Integrins mediate cell-matrix or cell-cell adhesion, and transduced signals that regulate gene expression and cell growth. This gene encodes the integrin beta 4 subunit, a receptor for the laminins. This subunit tends to associate with alpha 6 subunit and is likely to play a pivotal role in the biology of invasive carcinoma. Mutations in this gene are associated with epidermolysis bullosa with pyloric atresia. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

## **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 

Swiss Prot P16144

3691

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 28-310 of human Integrin beta 4 (ITGB4/CD104) (NP\_000204.3).

#### **Synonyms**

CD104; GP150; JEB5A; JEB5B; 4)

#### **Contact**

•

www.abclonal.com

### **Product Information**

**Source** Rabbit **Isotype** IgG


**Purification**Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of lysates from wild type (WT) and Integrin beta 4 (ITGB4/CD104) knockout (KO) HeLa cells, using [KO Validated] Integrin beta 4 (ITGB4/CD104) Rabbit pAb (A19890) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.