# **MPC1** Rabbit pAb

Catalog No.: A20195 2 Publications



# **Basic Information**

### **Observed MW**

12kDa

### **Calculated MW**

12kDa

### Category

Polyclonal Antibody

### **Applications**

WB,IP,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene is part of an MPC1/MPC2 heterodimer that is responsible for transporting pyruvate into mitochondria. The encoded protein is found in the inner mitochondrial membrane. Defects in this gene are a cause of mitochondrial pyruvate carrier deficiency. Several transcript variants, some protein coding and one non-protein coding, have been found for this gene.

# **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IP** 0.5μg-4μg antibody for

200μg-400μg extracts

of whole cells

**ELISA** Recommended starting

concentration is 1

μg/mL. Please optimize the concentration based on your specific

assay requirements.

# Contact

www.abclonal.com

## **Immunogen Information**

**Gene ID**Swiss Prot
51660
Q9Y5U8

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

MPYCD; BRP44L; CGI-129; SLC54A1; MPC1


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using MPC1 Rabbit pAb (A20195) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Immunoprecipitation analysis of 300 µg extracts of K-562 cells using 3 µg MPC1 antibody (A20195). Western blot was performed from the immunoprecipitate using MPC1 antibody (A20195) at a dilution of 1:1000.