ABclonal www.abclonal.com

TriMethyl-Histone H3-K36 Rabbit mAb

Catalog No.: A20379 Recombinant 6 Publications

Basic Information

Observed MW

17kDa

Calculated MW

15kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IHC-P,IP,ChIP,ChIP-seq,ELISA,DB

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC50050

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB 1:1000 - 1:6000

DB 1:500 - 1:1000

IHC-P 1:2000 - 1:8000

IP 0.5μg-4μg antibody for

200µg-400µg extracts of whole cells

......

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay

Contact

ELISA

www.abclonal.com

Immunogen Information

Gene ID	Swiss Prot
8290/8350	Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3.4; H3/g; H3FT; H3t; HIST3H3; Histone H3; HIST1H3A; TriMethyl-Histone H3-K36


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using TriMethyl-Histone H3-K36 Rabbit mAb (A20379) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.








Dot-blot analysis of all sorts of peptides using TriMethyl-Histone H3-K36 antibody (A20379) at 1:1000 dilution.

Immunohistochemistry analysis of paraffin-embedded Human colon tissue using TriMethyl-Histone H3-K36 Rabbit mAb (A20379) at a dilution of 1:5000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.


Immunohistochemistry analysis of paraffin-embedded Mouse liver tissue using TriMethyl-Histone H3-K36 Rabbit mAb (A20379) at a dilution of 1:5000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.









Immunohistochemistry analysis of paraffin-embedded Rat lung tissue using TriMethyl-Histone H3-K36 Rabbit mAb (A20379) at a dilution of 1:5000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunoprecipitation analysis of 600 µg extracts of 293F cells using 5 µg TriMethyl-Histone H3-K36 antibody (A20379). Western blot was performed from the immunoprecipitate using TriMethyl-Histone H3-K36 antibody (A20379) at a dilution of 1:1000.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using TriMethyl-Histone H3-K36 antibody (A20379) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Chromatin immunoprecipitations were performed with cross-linked chromatin from HeLa cells and H3K36me3 Rabbit mAb (A20379). The ChIP sequencing results indicate the enrichment pattern of H3K36me3 in selected genomic region and representative gene loci (GAPDH), as shown in figure.