

A20680

Leader in Biomolecular Solutions for Life Science



# MonoMethyl-Histone H3-K18 Rabbit mAb

Catalog No.: A20680

Recombinant

## Basic Information

### Observed MW

17kDa

### Calculated MW

16kDa

### Category

SMab Recombinant Monoclonal Antibody

### Applications

WB,IHC-  
P,IF/ICC,IP,ChIP,ELISA,DB,CUT&Tag

### Cross-Reactivity

Human,Mouse,Rat,Other (Wide Range Predicted)

### CloneNo number

ARC2621

## Recommended Dilutions

WB 1:500 - 1:1000

DB 1:500 - 1:1000

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

IP 0.5µg-4µg antibody for  
200µg-400µg extracts  
of whole cellsELISA Recommended starting  
concentration is 1  
µg/mL. Please optimize  
the concentration  
based on your specific  
assay requirements.ChIP 5µg antibody for  
5µg-10µg of ChromatinCUT&Tag 10<sup>5</sup> cells /1 µg

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

## Immunogen Information

### Gene ID

8290/8350

### Swiss Prot

Q16695/P68431

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; MonoMethyl-Histone H3-K18

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


Contact

---

 [www.abclonal.com](http://www.abclonal.com)

---

## Validation Data



Western blot analysis of various lysates using MonoMethyl-Histone H3-K18 Rabbit mAb (A20680) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.




Dot-blot analysis of all sorts of peptides using MonoMethyl-Histone H3-K18 antibody (A20680) at 1:1000 dilution.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer using MonoMethyl-Histone H3-K18 Rabbit mAb (A20680) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse lung using MonoMethyl-Histone H3-K18 Rabbit mAb (A20680) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain using MonoMethyl-Histone H3-K18 Rabbit mAb (A20680) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunofluorescence analysis of NIH/3T3 cells using MonoMethyl-Histone H3-K18 Rabbit mAb (A20680) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of U-2 OS cells using MonoMethyl-Histone H3-K18 Rabbit mAb (A20680) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.


Validation Data



Immunoprecipitation analysis of 600 µg extracts of 293F cells using 5 µg MonoMethyl-Histone H3-K18 antibody (A20680). Western blot was performed from the immunoprecipitate using MonoMethyl-Histone H3-K18 antibody (A20680) at a dilution of 1:1000.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using MonoMethyl-Histone H3-K18 antibody (A20680) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



CUT&Tag was performed using the CUT&Tag Assay Kit (pAG-Tn5) for Illumina(RK20265) from 10<sup>5</sup> K562 cells with 1µg MonoMethyl-Histone H3-K18 Rabbit mAb(A20680), along with a Goat Anti-Rabbit IgG(H+L). The CUT&Tag results indicate the enrichment pattern of H3K18Me1 in representative gene loci (MYT1), as shown in figure.