

A21128

Leader in Biomolecular Solutions for Life Science



## MLD Rabbit mAb

Catalog No.: A21128

Recombinant

### Basic Information

#### Observed MW

34kDa

#### Calculated MW

38kDa

#### Category

SMab Recombinant Monoclonal  
Antibody

#### Applications

WB,ELISA

#### Cross-Reactivity

Human

#### CloneNo number

ARC2977

### Background

This gene encodes a member of the membrane fatty acid desaturase family which is responsible for inserting double bonds into specific positions in fatty acids. This protein contains three His-containing consensus motifs that are characteristic of a group of membrane fatty acid desaturases. It is predicted to be a multiple membrane-spanning protein localized to the endoplasmic reticulum. Overexpression of this gene inhibited biosynthesis of the EGF receptor, suggesting a possible role of a fatty acid desaturase in regulating biosynthetic processing of the EGF receptor.

### Recommended Dilutions

WB 1:500 - 1:1000

### Immunogen Information

#### Gene ID

8560

#### Swiss Prot

O15121

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-100 of human MLD (NP\_003667.1).

#### Synonyms

MLD; DEGS; DES1; Des-1; FADS7; HLD18; MIG15; DEGS-1

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification

Affinity purification


#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## Validation Data

---



Western blot analysis of lysates from U-87MG cells, using MLD Rabbit mAb (A21128) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.