MLD Rabbit mAb

Catalog No.: A21128 Recombinant



Basic Information

Observed MW 34kDa

Calculated MW 38kDa

Category

SMab Recombinant Monoclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human

CloneNo number ARC2977

Recommended Dilutions

WB	1:500 - 1:1000
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Background

This gene encodes a member of the membrane fatty acid desaturase family which is responsible for inserting double bonds into specific positions in fatty acids. This protein contains three His-containing consensus motifs that are characteristic of a group of membrane fatty acid desaturases. It is predicted to be a multiple membrane-spanning protein localized to the endoplasmic reticulum. Overexpression of this gene inhibited biosynthesis of the EGF receptor, suggesting a possible role of a fatty acid desaturase in regulating biosynthetic processing of the EGF receptor.

Immunogen Information

Gene ID 8560

Swiss Prot 015121

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

MLD; DEGS; DES1; Des-1; FADS7; HLD18; MIG15; DEGS-1

Product Information

www.abclonal.com

Isotype IgG **Purification** Affinity purification


Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from U-87MG cells, using MLD Rabbit mAb (A21128) at1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.