Leader in Biomolecular Solutions for Life Science

# FT Rabbit pAb

Catalog No.: A21280



### **Basic Information**

### **Observed MW**

20kDa

### **Calculated MW**

20kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Arabidopsis thaliana

## **Background**

FT, together with LFY, promotes flowering and is antagonistic with its homologous gene, TERMINAL FLOWER1 (TFL1). FT is expressed in leaves and is induced by long day treatment. Either the FT mRNA or protein is translocated to the shoot apex where it induces its own expression. Recent data suggests that FT protein acts as a long-range signal. FT is a target of CO and acts upstream of SOC1.

### **Recommended Dilutions**

WB

1:500 - 1:1000

### **Immunogen Information**

**Gene ID** 842859

Swiss Prot Q9SXZ2

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 70-150 of arabidopsis thaliana FT (NP\_176726.1).

### **Synonyms**

F5I14.3; F5I14\_3; FLOWERING LOCUS T; REDUCED STEM BRANCHING 8; RSB8; FT

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of various tissues from Arabidopsis thaliana, using FT Rabbit pAb (A21280) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.