BES1 Rabbit pAb

Catalog No.: A21282



Basic Information

Observed MW

43kDa

Calculated MW

36kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Arabidopsis thaliana

Background

Encodes brassinosteroid (BR) signalling protein that accumulates in the nucleus as dephosphorylated form in response to BRs. Is phosphorylated by the BIN2 GSK3 kinase. It synergistically interacts with BIM1 to bind to E box sequences (CANNTG). The protein contains a nuclear localization signal (NLS), followed by a highly conserved aminoterminal domain (N) shared by all family members, a BIN2 phosphorylation domain (P), a PEST motif, involved in protein degradation in the absence of BR, and a carboxylterminal domain. BES1 can interact with the ELF6 and REF6 Jumonji N/C-domain containing proteins and may direct them to modify histone methylation upstream of some brassinosteroid responsive-genes. Works with BRAVO to regulate QC division in the root.

Recommended Dilutions

WB 1:500 - 1:1000

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID838518

Swiss Prot
Q9LN63

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

107 PROTEIN; BRASSINAZOLE-RESISTANT 2; BRI1-EMS-SUPPRESSOR 1; BZR2; F18O14.7; F18O14_7; BES1

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of extracts of various tissues from Arabidopsis thaliana, using BES1 Rabbit pAb (A21282) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.