A21702

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

[KO Validated] SQSTM1/p62 Rabbit pAb

Catalog No.: A21702 KO Validated 4 Publications

Basic Information

Observed MW
62kDa

Calculated MW
48kDa

Category
Polyclonal Antibody

Applications
WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity
Human,Mouse,Rat

Background

This gene encodes a multifunctional protein that binds ubiquitin and regulates activation of the nuclear factor kappa-B (NF-kB) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF-kB in response to upstream signals. Alternatively spliced transcript variants encoding either the same or different isoforms have been identified for this gene. Mutations in this gene result in sporadic and familial Paget disease of bone.

Recommended Dilutions

WB	1:2000 - 1:4000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

www.abclonal.com

Immunogen Information

Gene ID
8878

Swiss Prot
Q13501

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
p60; p62; A170; DMRV; OSIL; PDB3; ZIP3; p62B; NADGP; FTDALS3; 62

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃


Validation Data




Western blot analysis of various lysates using [KO Validated] SQSTM1/p62 Rabbit pAb (A21702) at 1:4000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 3s.




Western blot analysis of lysates from wild type(WT) and SQSTM1/p62 knockout (KO) 293T cells, using [KO Validated] SQSTM1/p62 Rabbit pAb (A21702) at 1:4000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 3s.



Immunohistochemistry analysis of paraffin-embedded Human lung adenocarcinoma using [KO Validated] SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human extranodal NK-T cell lymphoma using [KO Validated] SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney using [KO Validated] SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.


Validation Data




Immunohistochemistry analysis of paraffin-embedded Rat spleen using [KO Validated] SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunofluorescence analysis of NIH/3T3 cells using [KO Validated] SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of C6 cells using [KO Validated] SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:100. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of HeLa cells using SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of HepG2 cells using SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH/3T3 cells using SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue:



Immunofluorescence analysis of A-549 cells using SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue:




Immunofluorescence analysis of NIH/3T3 cells using SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue:

Validation Data

DAPI for nuclear staining.

DAPI for nuclear staining.

DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using SQSTM1/p62 Rabbit pAb (A21702) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.