

A21785

Leader in Biomolecular Solutions for Life Science



## [KD Validated] RARG Rabbit pAb

Catalog No.: A21785

### Basic Information

---

**Observed MW**

48kDa

**Calculated MW**

50kDa

**Category**

Polyclonal Antibody

**Applications**

WB,ELISA

**Cross-Reactivity**

Human

### Background

---

This gene encodes a retinoic acid receptor that belongs to the nuclear hormone receptor family. Retinoic acid receptors (RARs) act as ligand-dependent transcriptional regulators. When bound to ligands, RARs activate transcription by binding as heterodimers to the retinoic acid response elements (RARE) found in the promoter regions of the target genes. In their unbound form, RARs repress transcription of their target genes. RARs are involved in various biological processes, including limb bud development, skeletal growth, and matrix homeostasis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

### Recommended Dilutions

---

WB 1:500 - 1:1000

### Immunogen Information

---

**Gene ID**

5916

**Swiss Prot**

P13631

**Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-70 of human RARG (NP\_000957.1).

**Synonyms**

RARC; NR1B3; RARgamma; [KD Validated] RARG

### Contact

---



[www.abclonal.com](http://www.abclonal.com)

### Product Information

---

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of extracts of SK-BR-3 cells, using RARG antibody (A21785) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 90s.

Western blot analysis of extracts from wild type (WT) and RARG knockdown (KD) MCF7 cells, using RARG antibody (A21785) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25 $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 90s.

