MAP1LC3B Rabbit pAb

Catalog No.: A21800



Basic Information

Observed MW 14kDa,16kDa

Calculated MW 14kDa

Category **Polyclonal Antibody**

Applications WB,ELISA

Cross-Reactivity Human,Rat

Background

The product of this gene is a subunit of neuronal microtubule-associated MAP1A and MAP1B proteins, which are involved in microtubule assembly and important for neurogenesis. Studies on the rat homolog implicate a role for this gene in autophagy, a process that involves the bulk degradation of cytoplasmic component.

Recommended Dilutions

WB	1:500 - 1:1000
ELISA	Recommended starting concentration is 1
	µg/mL. Please optimize
	the concentration
	based on your specific
	assay requirements.

Immunogen Information

Gene ID **Swiss Prot** Q9GZQ8 Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-50 of human MAP1LC3B (NP_073729.1).

Synonyms

81631

LC3B; ATG8F; MAP1LC3B-a; MAP1A/1BLC3; 3B

Product Information

Ð www.abclonal.com

Source Rabbit


Isotype IgG

Purification Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from 293T cells using MAP1LC3B Rabbit pAb (A21800) at 1:1000 dilution. 293T cells were treated by Chloroquine (50 μ M) at 37°C for 20 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.



Western blot analysis of lysates from C6 cells using MAP1LC3B Rabbit pAb (A21800) at 1:1000 dilution. C6 cells were treated by Chloroquine (50 μ M) at 37°C for 20 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.