

A22079

Leader in Biomolecular Solutions for Life Science



MonoMethyl-Histone H3-K9 Rabbit mAb

Catalog No.: A22079

Recombinant

Basic Information

Observed MW

17kDa

Calculated MW

16kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB,IHC-P,IF/ICC,ELISA,DB

Cross-Reactivity

Human,Mouse,Rat,Other (Wide Range
Predicted)

CloneNo number

ARC54621

Recommended Dilutions

WB 1:1000 - 1:5000

DB 1:1000 - 1:5000

IHC-P 1:5000 - 1:20000

IF/ICC 1:50 - 1:200

ELISA Recommended starting
concentration is 1
μg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Contact



www.abclonal.com

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; MonoMethyl-Histone H3-K9

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Western blot analysis of various lysates using MonoMethyl-Histone H3-K9 Rabbit mAb (A22079) at 1:2000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.




Dot-blot analysis of all sorts of peptides using MonoMethyl-Histone H3-K9 antibody (A22079) at 1:2000 dilution.




Immunohistochemistry analysis of paraffin-embedded Mouse testis tissue using MonoMethyl-Histone H3-K9 Rabbit mAb (A22079) at a dilution of 1:5000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using MonoMethyl-Histone H3-K9 Rabbit mAb (A22079) at a dilution of 1:5000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat liver tissue using MonoMethyl-Histone H3-K9 Rabbit mAb (A22079) at a dilution of 1:5000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunofluorescence analysis of HeLa cells using MonoMethyl-Histone H3-K9 Rabbit mAb (A22079) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using MonoMethyl-Histone H3-K9 Rabbit mAb (A22079) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Validation Data



Immunofluorescence analysis of PC-12 cells using MonoMethyl-Histone H3-K9 Rabbit mAb (A22079) at dilution of 1:100(40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.