A22197

Leader in Biomolecular Solutions for Life Science

www.abclonal.com

CD34 Rabbit mAb

Catalog No.: A22197 Recombinant 1 Publications

Basic Information

Observed MW
120kDa

Calculated MW
41kDa

Category
Monoclonal Antibody

Applications
WB,IHC-P,ELISA

Cross-Reactivity
Human

Clone number
ARC55664

Background

The protein encoded by this gene may play a role in the attachment of stem cells to the bone marrow extracellular matrix or to stromal cells. This single-pass membrane protein is highly glycosylated and phosphorylated by protein kinase C. Two transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB	1:1000 - 1:2000
IHC-P	1:5000 - 1:20000
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
947

Swiss Prot
P28906

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
CD34; CD34 molecule; GIG3; MORT1

Contact

www.abclonal.com

Product Information


Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage
Store at -20℃


Validation Data




Western blot analysis of various lysates, using CD34 Rabbit mAb (A22197) at1:2000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:200000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using CD34 Rabbit mAb (A22197) at a dilution of 1:12000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using CD34 Rabbit mAb (A22197) at a dilution of 1:12000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human placenta tissue using CD34 Rabbit mAb (A22197) at a dilution of 1:12000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using CD34 Rabbit mAb (A22197) at a dilution of 1:12000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human liver cancer tissue using CD34 Rabbit mAb (A22197) at a dilution of 1:12000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using CD34 Rabbit mAb (A22197) at a dilution of 1:12000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.