

# [KO Validated] CDKN1A/p21 Rabbit PolymAb®

Catalog No.: A22460 KO Validated

## **Basic Information**

### **Observed MW**

21kDa

### **Calculated MW**

18kDa

#### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse

# **Background**

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or -cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Multiple alternatively spliced variants have been found for this gene.

# **Recommended Dilutions**

WB 1:10000 - 1:40000

**IF/ICC** 1:1000 - 1:4000

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

 Gene ID
 Swiss Prot

 1026/ 12575
 P38936/P39689

#### **Immunogen**

Recombinant protein of human/mouse CDKN1A/p21.

#### **Synonyms**

P21; CIP1; SDI1; WAF1; CAP20; CDKN1; MDA-6; p21CIP1

### **Contact**

www.abclonal.com

## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.