ABclonal www.abclonal.com

LAMP2 Rabbit mAb

Catalog No.: A22482 Recombinant 5 Publications

Basic Information

Observed MW

100-130kDa

Calculated MW

45kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human

CloneNo number

ARC54762

Background

The protein encoded by this gene is a member of a family of membrane glycoproteins. This glycoprotein provides selectins with carbohydrate ligands. It may play a role in tumor cell metastasis. It may also function in the protection, maintenance, and adhesion of the lysosome. Alternative splicing of this gene results in multiple transcript variants encoding distinct proteins.

Recommended Dilutions

WB 1:40000 - 1:160000

IHC-P 1:200 - 1:2000

IF/ICC 1:200 - 1:1000

ELISA Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

Contact

www.abclonal.com

Immunogen Information

Gene ID	Swiss Prot
3920	P13473

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

DND; LAMPB; CD107b; LAMP-2; LGP-96; LGP110; LAMP2


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.



Western blot analysis of various lysates, using LAMP2 Rabbit mAb (A22482) at1:40000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000

dilution.


Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Human kidney using LAMP2 Rabbit mAb (A22482) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of HeLa cells using LAMP2 Rabbit mAb (A22482) at dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.