

A22700

Leader in Biomolecular Solutions for Life Science



# Neurogranin (NRGN) Rabbit mAb

Catalog No.: A22700

Recombinant

## Basic Information

### Observed MW

15kDa

### Calculated MW

8kDa

### Category

SMab Recombinant Monoclonal Antibody

### Applications

WB,IHC-P,ELISA

### Cross-Reactivity

Mouse,Rat

### CloneNo number

ARC57532

## Background

Neurogranin (NRGN) is the human homolog of the neuron-specific rat RC3/neurogranin gene. This gene encodes a postsynaptic protein kinase substrate that binds calmodulin in the absence of calcium. The NRGN gene contains four exons and three introns. The exons 1 and 2 encode the protein and exons 3 and 4 contain untranslated sequences. It is suggested that the NRGN is a direct target for thyroid hormone in human brain, and that control of expression of this gene could underlay many of the consequences of hypothyroidism on mental states during development as well as in adult subjects.

## Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:500 - 1:1000

## Immunogen Information

<b>Gene ID</b>	<b>Swiss Prot</b>
4900	Q92686

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 2-78 of human Neurogranin (NRGN). (NP\_001119653.1).

### Synonyms

RC3; hng; Neurogranin (NRGN)

## Contact

 [www.abclonal.com](http://www.abclonal.com)

## Product Information


<b>Source</b>	<b>Isotype</b>	<b>Purification</b>
Rabbit	IgG	Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.


## Validation Data



Western blot analysis of various lysates, using Neurogranin(NRGN) Rabbit mAb (A22700) at 1 : 1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25ug per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 180s.



Western blot analysis of lysates from Rat spleen, using Neurogranin(NRGN) Rabbit mAb (A22700) at 1 : 1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25ug per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Enhanced Kit (RM00021).  
Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using Neurogranin (NRGN) Rabbit mAb (A22700) at dilution of 1:1000 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.