# **DSCR1 Rabbit mAb**

Catalog No.: A23086 Recombinant



### **Basic Information**

#### **Observed MW**

Refer to figures

#### **Calculated MW**

28kDa

#### **Category**

SMab Recombinant Monoclonal Antibody

### **Applications**

IHC-P,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC58928

# **Background**

The protein encoded by this gene interacts with calcineurin A and inhibits calcineurin-dependent signaling pathways, possibly affecting central nervous system development. This gene is located in the minimal candidate region for the Down syndrome phenotype, and is overexpressed in the brain of Down syndrome fetuses. Chronic overexpression of this gene may lead to neurofibrillary tangles such as those associated with Alzheimer disease. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

**IHC-P** 1:50 - 1:200

**IF/ICC** 1:50 - 1:200

# **Immunogen Information**

**Gene ID**Swiss Prot
1827
P53805

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 60-245 of human DSCR1. (NP\_004405.3).

#### **Synonyms**

CSP1; DSC1; RCN1; DSCR1; MCIP1; ADAPT78

#### **Contact**

www.abclonal.com

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification


## Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

## **Validation Data**







Immunohistochemistry analysis of paraffin-embedded Human kidney using DSCR1 Rabbit mAb (A23086) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Bufferr (pH 6.0) prior to IHC staining.

Immunofluorescence analysis of U-2 OS cells using DSCR1 Rabbit mAb (A23086) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of U-251MG cells using DSCR1 Rabbit mAb (A23086) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.